

## First Serological detection of *Leishmania* spp of domestic and stray cats in Al-Qadisiyah Province, Iraq

Kazem Najm Abdhullah AL-Husseini<sup>1</sup> Mansour Jadaan Ali<sup>2</sup>

<sup>1</sup>Al-Diwaniyah Veterinary Teaching Hospital, <sup>2</sup>Department of Microbiology, College of Veterinary Medicine, University of Al-Qadisiyah, Al-Diwaniyah City, Iraq

Submitted: December 29, 2024

Revised: January 28, 2025

Accepted: February 18, 2024

**Abstract** Leishmaniasis is a major mosquito-borne disease due to protozoa of *Leishmania* genus which spread by sand flies' bites. This study was conducted to detect Leishmaniasis in domestic and stray cats by using serological test (Indirect Enzyme-Linked Immunosorbent Assay -ELISA) and to determine the effects of sex, age and months on the infection rate in Al-Qadisiyah Province. A total of 142 samples were collected during the period from the first of July to the end of December/ 2024. Leishmaniasis IgG antibodies were detected rate 20.42% (29/142) of the cats by indirect ELISA. High infection rate was recorded in stray cats 28.37% (21/73), compared to domestic cats 11.76% (8/68), with significant difference at  $P<0.05$ . the prevalence rate was in males 23.28% (17/73) and in females 17.39% (12/69) without significant difference at  $P\geq0.05$ . Moreover, the highest infection rate in domestic cats was observed in the adult was 15.55% (7/45) while the lowest in kittens 4.34% (1/23), without significant difference at  $P<0.05$ . The prevalence was altered during various months studied where the highest and lowest rate was recorded (34.61%) and (4.54%) in September and July respectively, without significant difference at  $P\geq0.05$ . For the first time, Leishmaniasis was recorded to infected stray and domestic cats in Iraq

**Keywords:** Al-Qadisiyah, Blood, cats, ELISA, Leishmaniasis

©Authors, 2025, College of Veterinary Medicine, University of Al-Qadisyah. This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/> ).

**Introduction** *Leishmania* a zoonotic human disease, is hyperendemic in subtropical and tropical areas of the world. Visceral leishmaniasis (VL) influences approximate 12 million individuals worldwide, with an incidence rate ranging from 0.2-0.4% (1). *Leishmania* spp. are spread by a variety of female sandfly species in the Mediterranean basin, which bite the definitive hosts (vertebrates) during feeding time (2). Although cats were once thought to be less vulnerable to *L. infantum*, they are now understood to be both competent hosts for the parasite and a source of blood for its sand fly vectors (3,4). Moreover, feline leishmaniasis (FeL) attributed to *L. infantum* has been recorded worldwide, precisely in the Mediterranean areas (5,6,7).

Hepatomegaly, lymphadenopathy, ulcers, and eye lesions and nodules are the most common clinical signs in these cases (8). Granulomatous inflammation with many amastigotes has also been seen in mucosal nodules, kidney, liver, spleen, and the eyes (9,10). cutaneous, mucosal, and visceral signs are among the

disease's clinical manifestations. Systemic and high mortality participation of the organism was linked to visceral signs. Significant morbidity is often caused by the spread of parasites through other tissues, which is often linked to cutaneous or mucocutaneous symptoms (11). A comparative analysis of the complement system's activation in various hosts following stimulation with *L. infantum* revealed that cats were less susceptible to infection because their classical and alternative pathways were less activated, and their lectin pathway was more activated because of less C4b deposition (12).

*Leishmania* spp investigate in cats involved, *L. braziliensis*, *L. amazonensis*, *L. venezuelensis*, *L. mexicana* and *L. infantum* (13). various worldwide studies confirmed a high seroprevalence of FeL leishmaniasis was detected by IFAT and ELISA (14,15,16).

The reaction of antibodies presents in the sera with purified and soluble *Leishmania* antigens derived from in vitro culture is what drives ELISA (17). By

ELISA, the approach has been extremely useful for the investigation of anti-Leishmania antibodies.

#### Materials and methods

##### Ethical approve

This study was approved for the animal care at the College of Veterinary Medicine, University of Al-Qadisiyah, Under the No. 4955 dated 17/11/2024

##### Collection of samples

Using single-use syringes and needles, one hundred and forty-two blood samples were collected from the cephalic vein or jugular vein of stray and domestic cats randomly selected from the areas of Al-Qadisiyah Governorate from July to December 2024. And transferred to anticoagulant-free tubes. Then, the blood was centrifuged for 12 minutes at a speed of about 3000 rpm, and one ml of the serum was transferred to fine plastic tubes and saved at -20°C for serological examination. (19,20).

##### Indirect ELISA

In this study, an indirect ELISA test was used to detect IgG antibodies in the serum of the studied cats against leishmaniasis. The manufacturer's instructions for the ELISA test kit for the leishmaniasis were followed, and serum samples from all the studied cats were prepared with the kit materials and reagents. The optical density (O.D.) was measured using a microtiter plate reader (BioBase, China) at 450 nm within 20 min after analysis. The results were calculated using the generated standard curve by plotting the average O.D. originated for each of the standard concentrations on the horizontal (X) axis as well as the vertical (Y) axis concentration and the positive sensitivity result is 0.1 ng/ml (21,22).

##### Results

The total infection rate of feline leishmaniasis in blood samples examined were 20.42% (29/142). Table (1) shows a higher infection rate of Feline Leishmaniasis in stray cats 28.37% (21/74), more than domestic cats 11.76% (8/70). with significant variance.

**Table 1:** Total infection rate of Leishmaniasis according to type of cats.

Cats	No. of examined samples	Positive	Percentage (%)
Stray	74	21	28.37
Domestic	68	8	11.76
Total	142	29	20.42
Chi-Square( $\chi^2$ )	6.01		

P value		0.014 (HS)
---------	--	------------

HS: Highly significant difference at  $p<0.05$

##### Infection rate according to sex

A higher infection rate was detected in males Cat 23.28% (17/75), while the lowest in females Cat 17.39% (12/71), without significant difference  $P > 0.05$  (Table 2).

**Table 2:** Infection rate of Feline Leishmaniasis according to sex by using indirect ELISA.

Sex	No. of examined samples	Positive	Percentage (%)
Males	73	17	23.28
Females	69	12	17.39
Chi-Square ( $\chi^2$ )	0.759		
P value	0.384 (No)		

NS: No significant difference at  $p<0.05$

##### Infection rate in domestic Cats according to age

Adult cats, which showed a high infection prevalence than kitten cats were recorded a lower rate. There was significant variability (Table 3).

**Table 3:** Infection rate of Feline Leishmaniasis according to age by indirect ELISA

Age	No. of examined samples	Positive	Percentage (%)
Kittens	23	1	4.34
Adults	45	7	15.55
Total	68	8	11.76
Chi-Square ( $\chi^2$ )	1.84		
P value	0.175 (NS)		

NS: No significant difference at  $p<0.05$

##### Infection rate according to months

The infection rate of Feline Leishmaniasis was higher in September (34.61%), followed by October (29.62%), and there was low infection rate in December (9.52%), followed by July (4.54%) (Table 4).

**Table 4:** Infection rate of Feline Leishmaniasis according to months by indirect ELISA.

Months	No. of examined samples	Positive	Percentage (%)
July	22	1	4.54

August	22	5	22.72
September	26	9	34.61
October	27	8	29.62
November	24	4	16.66
December	21	2	9.52
Chi-Square ( $\chi^2$ )		9.85	
P value		0.079 (NS)	

NS: No significant difference at  $p<0.05$

### Discussion

Leishmaniasis is a complicated fly-borne disease originated by different species of Leishmania that is distributed in 80 countries of South and Central America, Asia and Africa, greater than 350 million individuals are calculated at risk of Leishmania (23). The objective of this study was investigation the existence of against Leishmania antibodies based on serological techniques in stray and domestic cats in Al-Qadisiyah Province Iraq. This study is the first detection of leishmaniasis in cats from Iraq where the incidence of leishmaniasis was recorded at 20.42%. And agreement with a study in Greece 23.2% (24), in Iran 24.27% (25), in Portugal 24.7% (26), in southern Italy 25.8% (27), in center Italy 16.3% (28), in Turkey 15.2% (29). and differed with the results of the study France 13.3% (30), in Italy 12.6% (31), in Jerusal 6.7% (32), in Pernambuco 3.9% (33), In Greece (Thessaloniki) 3.87% (34), 0% in Angola (35). 20.42% of examined cats were positive for antibodies against Leishmania using ELISA. The distribution of feline leishmaniasis was different between areas, and when comparing this investigation with other studies, the variables originated from various serological examinations and cross-reactions with other Leishmania species or may be due to different levels of distribution, types of feline populations involved in these studies, and validation of the serological techniques. (36).

The current study showed a higher infection rate of Feline Leishmaniasis in stray cats, 28.37% more than domestic cats, 11.76%. with found significant variance. And agreement with a study in the Mediterranean Basin is 11.9%, and domestic is 8%. recorded a high positive in sheltered cats in comparison with private-owned cats in the province of Paraná, BziParaná, in greater positivity that founded in sheltered cats may be due to their continuous exposure to infected mosquitoes, as the shelter examined in previous in a previous study surrounded

by open sewage and lack of basic sanitation as well as vegetation that supports the existence of vectors. Moreover, living together with infected dogs may pose the likelihood of feline leishmaniasis (37).

The current study showed the highest infection rate was recorded in males at 23.28%, while the lowest in females was 17.39% without significant difference. This study found agreement with the study in cats from southern Iran. Female 12.33% and male 38.61% (38). The results of our study differ from the study of Italy and Greece male 2.1% and female 3.9%. (39). For behavioral and environmental reasons, they are more prone to feline leishmaniasis. Male cats are more likely to roam around and fight with other cats, which raises the possibility that they will get bitten by disease-transmitting insects. According to some research, sex hormones might influence how the immune system reacts, making males less immune to infection than females. in adult cats showed a higher infection prevalence of 15.15% compared to kittens, which have a lower positivity rate of 4.34%. There was a large variation in the age of the cats, and nearly in agreement with a study conducted in the Mediterranean basin, the result was 13% in adult cats and 9.3% in kittens (40).

unexpectedly, adult cats have the higher risk of infection with *L. infantum* in comparison with other groups of age (41). this study the infection rate of Feline Leishmaniasis was higher September (34.61%), and lower rate in July (4.54%). was different from, the study in northern Italy where it was recorded higher in August 33.3 %, and there was low infection rate in October 5.6%, (42).

Temperature, precipitation, and humidity all have a significant impact on the behavior and distribution of sand flies (43). By altering the number of sand flies or the air temperature, which indicates the parasite's development and growth in the vector host, climate change and ambient temperature can have an impact on the spread of Leishmania. As for our current study, it is an increase in the prevalence in cats that were sampled in September and October, due to the increased activity of the *Phlebotomus* species in Al-Qadisiyah Province of Iraq during these two months.

### Conclusions

This study provides, for the first-time serological detection of Leishmania spp, in stray and domestic cats in Al-Qadisiyah Province, Iraq. Indicates that cats are regularly bitten by infected sand flies in and may

have a potential reservoir role in the maintenance of *L. spp.* moreover, additional studies are required to determine the role of cats in the distribution of this parasite and regardless of if this species should be actually considered closely reservoir of current parasite.

#### Acknowledgment

Special thanks and appreciations are paid to Head and all staff member of The Department of Microbiology, College of Veterinary Medicine, University of Al-Qadisiyah for the facilities they provided for me to conduct the research.

#### Conflict of interest

The authors declare that there is no conflict of interest of this paper.

#### Funding source

This research had no specific fund; however, it was self-funded by the authors.

#### References

1. Strauss-Ayali D, Jaffe CL, Burshtain O, Gonen L, Baneth G. Polymerase chain reaction using noninvasively obtained samples, for the detection of *Leishmania infantum* DNA in dogs. *J Infect Dis*, (2004);189(9):1729-33. DOI: 10.1086/383281
2. Tzani M, Barrasa A, Vakali A, Georgakopoulou T, Mellou K, Pervanidou D. Surveillance data for human Leishmaniasis indicate the need for a sustainable action plan for its management and control, Greece, 2004 to 2018, *Euro Surveill*, (2021);26(18):2000159. DOI: 10.2807/1560-7917.ES.2021.26.18.2000159
3. Asfaram S, Fakhar M, Teshnizi SH, Is the cat an important reservoir host for visceral leishmaniasis? A systematic review with meta-analysis. *J Venom Anim Toxins Incl Trop Dis*, (2019);25(1):e20190012. <https://doi.org/10.1590/1678-9199-JVATI TD-2019-0012>.
4. Nascimento LFJ, Cirilo TM, Gomes DS, Gomes ACA, Lima VFS, Scher R, Epidemiological and diagnostic aspects of feline leishmaniasis with emphasis on Brazil: a narrative review. *Parasitol Res*, (2022);121(1):21-34. DOI: 10.1007/s00436-021-07372-9
5. Spada E, Perego R, Vitale F, Bruno F, Castelli G, Tarantola G, Baggiani L, Magistrelli S, Proverbio D. Feline Leishmania spp. infection in a non-endemic area of northern Italy. *Animals (Basel)*, (2020); 10(5):817. DOI:10.3390/ani10050817
6. Alcover MM, Basurco A, Fernandez A, Riera C, Fisa R, Gonzalez A, Verde M, Garrido AM, Ruiz H, Yzuel A, Villanueva-Saz S. A cross-sectional study of *Leishmania infantum* infection in stray cats in the city of Zaragoza (Spain) using serology and PCR. *Parasit Vectors*, (2021); 14: 1-14. DOI:10.1186/s13071-021-04682-w
7. Muz, D., Can, H., Karakavuk, M., Döş, kaya, M., Ozdemir, H.G., Değirmenci Döş, kaya, A., Atalay, S., ahar, E., Pektaş, B., Karakus, M., Töz, S., Ozbel, Y., Gürüz, A.Y., Muz, M.N., The molecular and serological investigation of feline immunodeficiency virus and feline leukemia virus in stray cats of Western Turkey. *Comp. Immunol. Microbiol. Infect. Dis*, (2021); 78: 101688 DOI:10.1016/j.cimid.2021.101688
8. Abramo F, Albanese F, Gattuso S, Randone A, Fileccia I, Dedola C. Skin lesions in feline leishmaniasis: A systematic review. *Pathogens*, ( 2021);10(4):472. DOI: 10.3390 /pathogens10040472
9. Verneuil M. Ocular leishmaniasis in a cat: case report. *J Fr Ophthalmol* (2013); 36(4):e67-72. DOI: 10.1016/j.jfo.2012.09.006
10. Migliazzo A, Vitale F, Calderone S, Puleio R, Binanti D, Abramo F. Feline leishmaniasis: a case with a high parasitic burden. *Vet Dermatol*, (2014); 26(1): 69-70. DOI:10.1111/vde.12180
11. Gramiccia M. Recent advances in leishmaniasis in pet animals: epidemiology, diagnostics and anti-vectorial prophylaxis. *Vet Parasitol*, ( 2011); 181(1): 23-30. DOI: 10.1016/j.vetpar.2011.04.019
- 12- Tirado TC, Bavia L, Ambrosio AR, Campos MP, de Almeida SM, Messias Reason IJ, Figueiredo FB. A comparative approach on the activation of the three complement system pathways in different hosts of visceral leishmaniasis after stimulation with *Leishmania infantum*. *Developmental and Comparative Immunology*,(2021);120(1):104061. DOI: 10.1016/j.dci.2021.104061
13. Pennisi MG, Cardoso L, Baneth G, Bourdeau P, Koutinas A, Miró G, et al. LeishVet update and recommendations on feline leishmaniasis. *Parasit Vectors*, (2015); 8(1): 302. doi:10.1186/s13071-015- 0909-z. PMid:26041555
14. Vides JP, Schwartzt TF, Sobrinho LSV, Marinho M, Laurenti MD, Biondo AW, Leutenegger C, Marcondes M. *Leishmania chagasi* infection in cats with dermatologic lesions from an endemic area of visceral leishmaniasis in Brazil. *Vet Parasitol*, (2011); 178:22-28. DOI:10.1016/j.vetpar.2010.12.042
15. Sobrinho LSV, Rossi CN, Vides JP, Braga ET, Gomes AAD, Lima VMF, Perri SHV, Generoso D, Langoni H, Leutenegger C, Biondo AW, Laurenti MD, Marcondes M. Coinfection of *Leishmania chagasi* with *Toxoplasma gondii*, feline immunodeficiency virus (FIV) and feline leukemia virus (FeLV) in cats from an endemic area of zoonotic visceral leishmaniasis. *Vet Parasitol*, (2012); 187(1-2):302-306. DOI:10.1016/j.vetpar.2012.01.010
16. Matos AMRN, Caldart ET, Ferreira FP, Monteiro KC, Souza M, Brunieri DTSC, Hilst CLS, Mascarenhas NMF, Mitsuka-Breganó R, Freire RL, Navarro IT. Antibodies anti-trypanosomatides in domestic cats in Paraná: who is at highest risk of infection? *Rev Bras Parasitol Vet*, (2018); 27(1):232-236. DOI: org/10.1590/s1984-296120180033
17. Silveira Neto L, Marcondes M, Bilsland El. Clinical and epidemiological aspects of feline leishmaniasis in Brazil.

Semina: Ciências Agrárias, Londrina, (2015);36(3):1467-1480. DOI: 10.5433/1679-0359.2015v36n3p1467

18. Trevisan DAC, Lonardoni MVC, Demarchi IG. Diagnostic methods to cutaneous leishmaniasis detection in domestic dogs and cats. *An Bras Dermatol*, (2015);90(6):868-72. DOI: 10.1590/abrd1806-4841.20153716

19. Carbonara M, Iatta R, Miró G, Montoya A, Benelli G, Mendoza-Roldan JA, Papadopoulos E, Lima C, Bouhsira E, Nachum-Biala Y, Decaro N, Schunack B, Baneth G, Otranto D. Feline leishmaniosis in the Mediterranean Basin: a multicenter study. *Parasites & Vectors*, (2024); 17:346. DOI: 10.1186/s13071-024-06419-x

20. Alcover MM, Basurco A, Fernandez A, Riera C, Fisa R, Gonzalez A, Verde M, Garrido AM, Ruíz H, Yzuel A, Villanueva-Saz S. A cross-sectional study of Leishmania infantum infection in stray cats in the city of Zaragoza (Spain) using serology and PCR. *Parasites Vectors*, (2021); 14(1):178. DOI: 10.1186/s13071-021-04682-w

21. Pradella GD, Escobar TA, Santos TP, Minuzzi JS, Silva LKR, Roman IJ, Vogel FSF, Duarte CA, Lübeck I. Detection of antibodies against Leishmania species using enzyme-linked immunosorbent assay in cats from the western border of Brazil. *Ciência Rural*, (2024); 54(1):8. DOI: 10.1590/0103-8478cr20230127

22. Médici KC, Navarro IT, Mitsuka-Breganó R, Caldart E. Seroprevalence of feline leishmaniasis in Paraná using antigens from different species of Leishmania spp. *Brazilian Journal of Development*, (2021);7(3):. 2525-8761. DOI: 10.34117/bjdv7n3-562

23. Abramo F, Albanese F, Gattuso S, Randone A, Fileccia I, Dedola C, Ibba F, Ottaiano P, Brianti E. Skin lesions in feline leishmaniosis: a systematic review. *Pathogens*, (2021) 10(4):472. DOI: 10.3390/pathogens10040472

24. Baneth G, Nachum-Biala Y, Zuberi A, Zipori-Barki N, Orshan L, Kleinerman G. Leishmania infection in cats and dogs housed together in an animal shelter reveals a higher parasite load in infected dogs despite a greater seroprevalence among cats. *Parasit Vectors*, (2020);13(1):115. DOI: 10.1186/s13071-020-3989-3

25. Mohebali M, Malmasi A, Khodabakhsh M, Zarei Z, Akhouni B, Hajjaran H. Feline leishmaniosis due to Leishmania infantum in Northwest Iran: the role of cats in endemic areas of visceral leishmaniosis. *Vet Parasitol Reg Stud Rep*, (2017); 9(1): 13-16. DOI: 10.1016/2017.03.010

26. Symeonidou I, Sioutas G, Gelasakis AI, Tsokana CN, Papadopoulos E. Leishmaniosis in Greece: the veterinary perspective. *Pathogens*, (2023);12(6):769. DOI: 10.3390/pathogens12060769

27. Otranto D, Napoli E, Latrofa MS, Annoscia G, Tarallo VD, Greco G. Feline and canine leishmaniosis and other vector-borne diseases in the Aeolian Islands: Pathogen and vector circulation in a confined environment. *Vet Parasitol*, (2017); 236(1):144-51. DOI: 10.1016/2019 PMID: 28288759

28. Vita S, Santori D, Aguzzi I, Petrotta E, Luciani A. Feline leishmaniasis and ehrlichiosis: serological investigation in Abruzzo region. *Vet Res Commun*, (2005); 2(1):319-21. DOI: 10.1007/s11259-005-007

29. Can H, Doğukan M, Özdemir HG, Şahar EA, Karakavuk M, Pektaş B. Seroprevalence of Leishmania infection and molecular detection of Leishmania tropica and Leishmania infantum in stray cats of İzmir, Turkey. *Exp Parasitol*, (2016); 167(1):109-14. DOI: 10.1016/j.exppara.2016.05.011

30. Galvez R, Montoya A, Cruz I, Fernandez C, Martin O, Checa R, et al. Latest trends in Leishmania infantum infection in dogs in Spain, Part I: mapped seroprevalence and sand fly distributions. *Parasit Vectors*, (2020);13(1):204. DOI: 10.1186/s13071-020-04081-7

31. Mendoza-Roldan J, Benelli G, Panarese R, Iatta R, Furlanello T, Beugne F, et al. Leishmania infantum and *Dirofilaria immitis* infections in Italy, 2009–2019: changing distribution patterns. *Parasit Vectors*, (2020);13(1):193. DOI: 10.1186/s13071-020-04063-9.

32. Nasereddin A, Salant H, Abdeen Z. Feline leishmaniasis in Jerusalem: serological investigation. *Vet Parasitol*, (2008);158(4):364-369. DOI: 10.1016/j.vetpar.2008.09.022

33. Silva RCN, Ramos RAN, Pimentel DS, Oliveira GMA, Carvalho GA, Santana MA. Detection of antibodies against Leishmania infantum in cats (*Felis catus*) from the state of Pernambuco, Brazil. *Rev Soc Bras Med Trop*, (2014); 47(1): 108-109. DOI: 10.1590/0037-8682-0005-2012

34. Chatzis MK, Andreadou M, Leontides L, Kasabalis D, Mylonakis M, Koutinas AF. Cytological and molecular detection of Leishmania infantum in different tissues of clinically normal and sick cats. *Vet Parasitol*, (2014); 202(3-4): 217-225. DOI: 10.1016/j.vetpar.2014.02.044. PMID: 24629427.

35. Lopes AP, Oliveira AC, Granada S, Rodrigues FT, Papadopoulos E, Schallig H. Antibodies to *Toxoplasma gondii* and Leishmania spp. in domestic cats from Luanda, Angola. *Vet Parasitol*, (2017); 239: 15-18. DOI: 10.1016/j.vetpar.2017.04.009

36. Nasereddin A, Salant H, Abdeen Z. Feline leishmaniasis in Jerusalem: serological investigation. *Vet Parasitol*, (2008); 158(4): 364-369. DOI: 10.1016/j.vetpar.2008.09.022. PMID: 18986768

37. Baneth G, Nachum-Biala Y, Zuberi A, Zipori-Barki N, Orshan L, Kleinerman G, Shmueli-Goldin A, Bellaiche M, Leszkowicz-Mazuz M, Salant H, Yasur-Landau D. Leishmania infection in cats and dogs housed together in an animal shelter reveals a higher parasite load in infected dogs despite a greater seroprevalence among cats. *Parasit Vectors*, (2020);13(1): 4-11. DOI: 10.1186/s13071-020-3989-3

38. Qasem A, Iraj M, Farzaneh B, Mohammad H M. Alarming: high prevalence of Leishmania infantum infection in cats from southern Iran based on molecular and serological methods. *Annals of Parasitology*, (2020);66(2):143-156. DOI: 10.17420/ap6602.249

39. Simone M, Mariasole C, Dimitris D, Alessandra B. Leishmania infantum Seroprevalence in Cats From Touristic

Areas of Italy and Greece. *Frontiers in Veterinary Science*, (2020); 11(7):616566 DOI: 10.3389/fvets.2020.616566

40. Mariaelisa C, Roberta I, Guadalupe M, Ana M, Giovanni B . Feline leishmaniosis in the Mediterranean Basin: a multicenter study. *Parasites & Vectors*, (2024);17(1):346. DOI:10.1186/s13071-024-06419-x

41. Atta R, Furlanello T, Colella V, Tarallo VD, Latrofa MS, Brianti E . A nationwide survey of *Leishmania infantum* infection in cats and associated risk factors in Italy. *PLoS Negl Trop Dis*, (2019) ; 13(7):e0007594. DOI: 10.1371/journal.pntd.0007594

42. Urbani L, Tirolo A, Salvatore D, (2020). Serological, molecular and clinicopathological findings associated with *Leishmania infantum* infection in cats in Northern Italy. *Feline Medicine and Surgery. and Surgery*. 22(10), 935-943. DOI:10.1177/1098612X19895067 Ready P, (2008). "Leishmaniasis emergence and climate change." *Rev SciTech*. 27(2), 399-412.

43. Carvalho, BM, Rangel EF, Ready PD, Vale MM. .Ecological niche modelling predicts southward expansion of *Lutzomyia (Nyssomyia) flaviscutellata* (Diptera: Psychodidae: Phlebotominae), vector of *Leishmania Leishmania amazonensis* in South America, under climate change. *PLoS One* ,(2015);10(11):e0143282.DOI:10.1371/journal.po