



The prevalence of methicillin-resistant *Staphylococcus aureus* in poultry farms

R.F. Mero and Dh.M. Jwher

Ramadhan Fattah Miro

Dep. of Vet. Public Health, College of Vet. Med. University of Mosul, Mosul, Iraq

*Corresponding author

ORCID: <https://orcid.org/0000-0002-7192-621X>

Email: diaataher@uomosul.edu.iq

©Authors, 2025, College of Veterinary Medicine, University of Al-Qadisiyah. This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

Submitted: February 02, 2025

Revised: March 10, 2025

Accepted: March 26, 2025

Abstract

The current study aimed to isolate and identify methicillin-resistant *Staphylococcus aureus* (MRSA) from poultry and study its prevalence in farms in northern Iraq. Two hundred and thirty-four samples were collected from broiler farms and their environment. The samples included swabs from chicken's skin, workers' hands, ventilators, feeders, drinking water, chicken feed, bedding, soil and grass, during the period from September 2024 till December 2024.

All samples were subjected to traditional and molecular methods for detection and identification for *S. aureus* and MRSA. The results showed that 43.16% of the total samples were positive for *S. aureus*. The highest isolation rate was from workers' hands and chickens at 53.85% from the total isolates, the lowest isolation rate was from feed and water at 34.61% depending on nuc gene. The results of MRSA isolation showed that 16.24% were positive for chromogenic agar and molecular methods depending on mecA gene, the highest rate was recorded from Bedding and Chicken at 34.61% and 26.92% respectively, and the lowest percentage of isolation was from Grass, Soil and Chicken feed at 3.85% and 7.69% respectively. The presence of MRSA emphasizes the need to implement continuous monitoring and surveillance programs and studies and collect data from other environments of livestock of all kinds, society and hospitals in order to develop insights into the relationship between human and animal strains, which is important for controlling and combating the microbe.

Keyword: *Staphylococcus aureus*, methicillin-resistant *Staphylococcus aureus*, poultry farms, Iraq

Introduction

Staphylococcus aureus (*S. aureus*) is one of the most important pathogens that threaten human life due to the high rates of infection and mortality it causes, despite its symbiotic presence with living organisms (1,2). The bacterium is highly tolerant to physicochemical environmental conditions, as it can survive in the open air for several days to weeks, Spreading for long distances, and is tolerant of drought (3,4). It also has the ability to remain alive for several weeks in dry pus (5). It can withstand a temperature of (60) °C for (30) minutes, but it dies after (60) minutes at the same temperature (6).

It's found in the surrounding environment such as water, soil and air. It has been isolated from foods of animal origin, and is a major cause of many cases of food poisoning around the world (14,15).

The investigation of the risk of human infection following contact with animals or their products, thereof, is still neglected in Iraq. The widespread, unrestricted use of antimicrobial compounds in food-producing animals has led to the emergence of MDR, making control and eradication of MRSA difficult. MRSA strains have been isolated from several types of food products, Including poultry and its products (16).

The most important sources of contamination in poultry farms with microbes are: workers, human waste, drinking water, feed, tools used in the field, rodents and hatcheries (17,18), In addition to the environment in which poultry are raised, such as water, soil, bedding, feces, waste, sick and dead birds, eggs, and other poultry products (19,20,21,22,23).

Transmission occurs in several ways, including inhalation of air, consumption of contaminated water and food, direct contact through hands or contact with secretions or contaminated materials and vectors (24,25), several studies conducted on poultry farms have revealed the presence of *S. aureus* in samples collected from humans, chickens, rodents, poultry litter, and soil surrounding the farm (26,27,28).

S. aureus have been associated with a number of conditions including dermatitis, omphalitis, femoral head necrosis, arthritis, tendinitis, and Bumble foot (26,29,30), many studies conducted on poultry, including live chickens, sick chickens and dead chickens, have proven the presence of *S. aureus* in large quantities (28,31).

From the above and the lack of studies that evaluate the extent of contamination with methicillin-resistant *S. aureus*, the current study aimed to isolate and identify MRSA from poultry and study its prevalence in farms in northern Iraq.

Material and Methods

Ethical approval

The approval for conducting the research was obtained from the Institutional Animal Care and Use Committee at the College of Veterinary Medicine at the University of Mosul, No. UM.VET.2024.046, in 9/7/2024.

Study area:

This study included 26 poultry farms from different regions and distributed in three governorates in the northern part of Iraq (Duhok, Nineveh and Erbil) selected randomly, during the period from September 2024 till December 2024.

Sampling:

Two hundred and thirty-four samples were collected from broiler farms and their environment. The samples included cotton swabs from chicken's skin, workers' hands, ventilators, feeders, drinking water, chicken feed, bedding, soil and grass, (26 samples each) placed in sterile tubes or containers containing peptone water. The samples were then delivered immediately to the Scientific Research Laboratory at the College of Veterinary Medicine / University of Mosul for bacteriological analysis.

Isolation of *S. aureus*

Microbiological methods described by Markey et al., (32), were followed to isolate *S. aureus* from the different samples, including cultivation on selective Mannitol Salt Agar (MSA) medium prepared by (Himedia®/India), Gram staining and cultivation on 5% sheep blood agar were used to test the hemolytic activity of the isolates, in addition to coagulase and catalase tests. CHROMagar™ (Himedia/India) was also used to identify MRSA (9).

Molecular Identification of *S. aureus*

Isolation of DNA:

Accurately following microbiological testing, the DNA of *S. aureus* isolates were extracted and analyzed. The samples were cultivated on MSA and incubated at 37°C for 24 hours. Then DNA was extracted from *S. aureus* isolates using Qiagen® (Germany) DNeasy Blood and Tissue Kit, according to the instructions. The concentration of extracted DNA was then measured with the Genova Nano (Jenway®/UK) instrument, and properly kept at -20°C.

Polymerase Chain Reaction technique (PCR)

As shown in Table (1), PCR technique was utilized to amplify particular sequences of the *nuc*, gene for *S. aureus* and *mecA* gene for MRSA isolates. A total of 25 µl was used for the PCR reaction mixture containing 12.5 µl of Promega Corporation's (2×) GoTaq Green Mix Master, 1 µl of the forward primer, 1µl of the reverse primer, 6.5 µl of Qiagen® (Germany) DNeasy-free water, and 4 µl of extracted DNA template made up the reaction mixture. The entire mixture was placed in a PCR tube, and the total volume was adjusted to 25 µl. The PCR amplification was performed under specific thermal cycling conditions Table (2). These conditions, including denaturation, annealing, and extension temperatures and durations, were tailored to the PCR protocol being used and optimized for the primer set and DNA template under the study. Next, 2% agarose gel electrophoresis Peqlab (Erlangen®/ Germany) was used to visualize the target sequence amplicons. A gel from along with a 100-3000 bp ladder DNA marker. Electrophoresis was carried out to separate and visualize the amplified DNA fragments, which were then compared to the DNA ladder for size estimation Table 1: Utilizing primers for identification of *S. aureus* and MRSA (*mecA* gene)

Gene	Primer	sequence	size (bp)	Reference
<i>nuc</i>	<i>nuc-F</i>	GCGATTGATGGTGATACGGTT	279	(33)
	<i>nuc-R</i>	AGCCAAGCCTTGACGAACTAAAGC		
<i>mecA</i>	<i>mecA-F</i>	GTGAAGATATACCAAGTGATT	147	(33)
	<i>mecA-R</i>	ATGCGCTATAGATTGAAAGGAT		

Table 2: PCR reaction program for *nuc* and *mec A* genes:

N	Steps	Temperature °C	Time min	Number of cycles
1	Initial denaturation	95	10	1
2	Denaturation	95	1	35
3	Annealing	60 (<i>nuc</i>) and 52 (<i>mecA</i>)	1	
4	Extension	72	1	
5	Final Extension	72	5	1

Results

The results of the initial isolation of *S. aureus* on MSA showed the growth of medium to large, round, smooth, soft, elevated and shiny yellow colonies with a change in the color of the medium from red to yellow. On blood agar, which caused beta hemolysis. Microscopically, the bacterial colonies were Gram-positive and grape-shaped, arranged in single cells or pairs, or forming irregular coccoid groups resembling grape clusters. On chromogenic medium, the bacterial colonies appeared bluish-green in color.

The isolation rates showed conformity between conventional and molecular techniques. Out of 234 samples examined, 101 samples were positive for isolation, i.e. 43.16% of the total samples included in the study (Workers hand, Chicken, Ventilator, Feeder and Water, Soil, Bedding, Grass and Chicken feed). The highest isolation rate was from workers' hands and chickens at 53.84%, and the lowest isolation rate was from feed and water at 34.61% using molecular techniques after electrophoresis of the amplification products of the nuc gene on agar gel, which were identical to the ladder size at 279 bp (Table 3 and Figure 1).

Table 3: Number and percentage of *S. aureus* isolates from different samples

n	Type of sample	No. of Samples	No. of +Ve	%
1	Workers hand	26	14	53.84
2	Chicken	26	14	53.84
3	Ventilator	26	12	46.15
4	Feeder	26	9	34.61
5	Water	26	9	34.61
6	Soil	26	12	46.15
7	Bedding	26	10	38.46
8	Grass	26	11	42.30
9	Chicken feed	26	10	38.46
Total		234	101	43.16




Fig. 1: PCR amplification product for *S. aureus* isolates for nuc gene at 279bp (M: 100-3000 bp ladder; 1-16: positive sample; 17: negative samples).

The highest prevalence rate of *S. aureus* was recorded among the samples, from the workers hands and chickens at 5.98%, while the lowest prevalence rate was in the feeder and water samples at 3.85%, as shown in figure (2).




Fig. 2: Prevalence of *S. aureus* of different studied samples.

The results of MRSA isolation showed that 38 samples out of 234 samples were positive, i.e. 16.23% by chromogenic agar medium and molecular methods by electrophoresis of amplification products for mecA gene at 147 bp on an agarose gel. The highest percentage of MRSA isolation was recorded from Bedding and Chicken at 34.61% and 26.92% respectively, and the lowest percentage of isolation was from Grass and Chicken feed at 3.84% and 7.69% respectively (Table 4 and Figure 3 and 4).

Table (4): Number and percentage of MRSA isolates from different samples

No.	Type of sample	No. of Samples	No. of +Ve	%
1	Workers hand	26	5	19.23
2	Chicken	26	7	26.92
3	Ventilator	26	3	11.53
4	Feeder	26	5	19.23
5	Water	26	4	15.38
6	Soil	26	2	7.69
7	Bedding	26	9	34.61
8	Grass	26	1	3.84
9	Chicken feed	26	2	7.69
Total		234	38	16.23




Fig. 3: Growth of methicillin-resistant *S.aureus* (MRSA) on MRSA chromogenic medium.




Fig. 4: PCR reaction product of MRSA isolates for the *mecA* gene at 147 bp (M: 100-3000 bp ladder; 1-8,11,13,16: positive sample;9,10,12,14,15,17: negative samples; 18: control negative)

The highest prevalence rate of MRSA was recorded among the samples of different fields and from different areas, from the bedding and chicken at 23.68% and 18.42% respectively, while the lowest prevalence rate was in the Grass, Soil and Chicken feed samples at 2.63%, 5.27% and 5.27% respectively, as shown in figure (figure 5).




Figure (5): Prevalence of MRSA of different studied samples

Discussion

Our study aimed to evaluate the level of *S. aureus* (MRSA) contamination, which is an indicator of the level of management and biosecurity measures followed in poultry farms, which have recently begun to grow and increase in Iraq in general.

In the northern regions of Iraq. Our study revealed a high prevalence of *S. aureus* and MRSA isolated from chickens, workers' hands, ventilators, feeders, drinking water, chicken feed, bedding, soil and grass.

Our study showed that out of a total of 234 samples, 101 samples, or 43.16%, were positive for *Staphylococcus aureus*, of which 38 samples, or 16.23%, were MRSA from all samples included in the study. Compared to studies conducted on workers and poultry, the study conducted by Assafi et al., (27) in Duhok province, recorded an isolation rate of 24% and

27.3%, respectively, which is higher than the rates we reached in our current study.

In comparison with studies conducted locally by (AL-Salihi et al., (34), a study conducted in Kirkuk Governorate on poultry workers recorded a rate of 16.6%, which is less than the rate we reached, which was 19.23%. The reason for the difference is likely to be due to the difference in the sample collection area, as the first was from the nose and the second was from the hands. Our results were completely similar to the results reached by (16) in Duhok Governorate, which reached 27.77% when compared to our results, which reached 26.92%. While the isolation results of MRSA from poultry meat varied between 40%, 33.3% and 14.81% in Iran, Wasit and Basra (13,35,36).

Internationally, studies have recorded varying isolation rates of MRSA from poultry including 71.5% in Germany (37). In contrast, the results of the study conducted in Egypt 36% (12), in Iran 67% (2), in addition to recording high levels of MRSA contamination in Austria, Denmark 52% (38) and 43.3% in Korea (39).

We did not observe any studies that addressed the isolation and diagnosis of MRSA from Ventilator, Feeder, Water, Soil, Bedding, Grass and Chicken feed in poultry farms, despite their importance in the occurrence of transmission and contamination (40), However, in an epidemiological study conducted by Hussein et al., (41) to investigate the presence of bacteria in the air inside poultry farms, it was revealed at a rate of 50%, which is a major cause of water and food contamination for workers, etc.

The variation in the isolation rates obtained in our study compared to other studies may be explained by many factors, including the administrative and health practices of poultry farms, breeding methods, the diagnostic methods used, geographical differences, etc., in addition to other factors like the environment in which poultry are raised, such as water, soil and bedding, feces and waste, sick and dead birds, and other poultry products, this is what is confirmed by Khan et al., (22) and Laban et al., (23).

The possession of the *mecA* gene by *S. aureus*, which adds to the bacteria another new mechanism for resistance to antibiotics, especially methicillin, is a criterion for diagnosing MRSA molecularly *S. aureus* isolates, as it is considered an inevitable criterion for confirming MRSA (42).

The results of the study were based on diagnosis and characterization by molecular methods in order to obtain the best and most accurate and sensitive results. The detection and diagnosis of *S. aureus* and MRSA

bacteria depends on traditional and molecular methods (43).

Direct contact by poultry farm workers with birds during field management operations is an important factor in the transmission of *S. aureus* from poultry to farm workers and vice versa (27), and therefore, *S. aureus* isolated from poultry is considered a risk indicator for humans living near poultry farms or dealing with them or with their production chains, as confirmed by the researcher (44,45). This may also occur due to the failure of biosecurity procedures, which include cleaning and sterilization of field components from chicken remains, or through handling birds for therapeutic purposes (41).

Therefore, reverse transmission from workers can also cause contamination and transmission of *Staphylococcus aureus* to birds and their environment if biosecurity procedures are not applied properly (8,9,27,41,45).

Conclusion

MRSA was isolated from broiler chickens, and its spread in the environment surrounding poultry houses is striking, as it constitutes a dangerous factor and indicator that threatens public health. The results highlight the importance of implementing biosecurity programs and adopting the Hazard Analysis and Critical Control Points (HACCP) system in poultry farms, placing restrictions and monitoring on the use of antibiotics, and applying strict health measures during poultry farming, as these measures are essential to prevent the spread of resistant bacteria and protect public health. Confirming the presence of MRSA emphasizes the need to implement continuous monitoring and surveillance programs and studies and collect data from other environments of livestock of all kinds, society and hospitals in order to develop insights into the relationship between human and animal strains, which is important for controlling and combating the microbe

Acknowledgment

The researchers would like to thank and appreciate the Deanship of the College of Veterinary Medicine / University of Mosul for its support and provision of the research environment. Thanks are also extended to those who contributed to the completion of this work by facilitating the collection of data.

Conflict of interest

The researcher declares that there is no conflict of interest regarding this work.

Funding source

This research had no specific fund; however, it was

self-funded by the authors.

References:

- 1- Pollitt E, Szkuta PT, Burns N, Foster SJ. *Staphylococcus aureus* infection dynamics. *PLoS Pathogens*.2018;14(6):211-215. <https://doi.org/10.1371/journal.ppat.1007112>
- 2- Rahimi F, Karimi S. Characteristics of Methicillin Resistant *Staphylococcus aureus* Strains Isolated From Poultry in Iran. *Archives of Clinical Infectious Diseases*.2015;10(4).<https://doi.org/10.5812/archcid.30885>
- 3- Carter GR, Cole JR. *Diagnostic Procedures in Veterinary Bacteriology and Microbiology*. 5th ed., Academic Press, Inc., Harcourt Brace Jovanovich Publishers, London, New York. 1990; PP. 201-207.
- 4- Kloos WE, Bannerman TL. *Staphylococcus and Micrococcus*. In: *Manual of clinical Microbiology* . Balows, A.; Hausler, W. J.; Herrmann, K. L.; Isenberg, H. D. and Shadomy, H. J.(Eds). 6th ed. American Society for Microbiology, Washington.1995.
- 5- Edmond MB, Wenzel RP, Pascalle AW. Vancomycin-Resistant *Staphylococcus aureus*: Perspectives on measures needed for control. *Annual of Internal Medicine* .1996;124(1):329-334.<https://doi.org/10.7326/0003-4819-124-3-199602010-00008>
- 6- Koneman EW, Allan SD, Janda WM. *Color Atlas and Textbook of Diagnostic Microbiology*, 5th ed. J.B. Lippincott, Philadelphia,1997.
- 7- Capurro A, Aspán A, Unnerstad EH, Waller PK, Artursson K. Identification of potential sources of *Staphylococcus aureus* in herds with mastitis problems. *Journal of Dairy Science*. 2010; 93(1):5-14. <https://doi.org/10.3168/jds.2009-2471>
- 8- Geenen PL, Graat EAM, Haenen A, Hengeveld PD, Van hoek AHAM, Huijsdens X W, Kappert CC, Lammers GAC, Van dujkeren E, Van DE Giessen AW. Prevalence of livestock-associated MRSA on Dutch broiler farms and in people living and/or working on these farms. *Epidemiology and Infection*. 2012; 141(5), 1099–1108. <https://doi.org/10.1017/s0950268812001616>
- 9- Rahma H, Jwher DH. Detection of *clfA*, *clfB* and *coa* genes in Methicillin-Resistant *Staphylococcus aureus* (MRSA) isolated from Nasal Cavity of Cows, Buffalo and their Breeders in Nineveh Governorate, Iraq. *Journal of Applied Veterinary Sciences*, 2024;9(3):1-10. <https://doi.org/10.21608/javs.2024.276537.1324>
- 10- Pal M, Kerorsa GB, Marami LM, Kandi V. Epidemiology, Pathogenicity, Animal Infections, Antibiotic Resistance, Public Health Significance, and



- Economic Impact of Staphylococcus Aureus: A Comprehensive Review. *American Journal of Public Health*. 2020; 8(1), pp.14-21. <https://www.sciepub.com/ajphr/abstract/11420>
- 11- Moosavian M, Shahin M, Navidifar T, Torabipour M. Typing of staphylococcal cassette chromosome mec encoding methicillin resistance in Staphylococcus aureus isolates in Ahvaz, Iran. *New Microbes and New Infections*. 2018; 21(1): 90–94. <https://doi.org/10.1016/j.nmni.2017.11.006>
- 12- Karmi M. Prevalence of methicillin-resistant Staphylococcus aureus in poultry meat in Qena, Egypt. *Veterinary World*. 2013;6(10):711–715. <https://doi.org/10.14202/vetworld.2013.711-715>
- 13- Jafarzadeh M, Meamar N, Mohsenzadeh M, Razmyar J. Identification of Methicillin-Resistant Staphylococcus aureus in poultry meat portions using Multiplex PCR. *Journal of Poultry Sciences and Avian Diseases*.2023;1(4):13–19. <https://doi.org/10.61838/kman.jpsad.1.4.3>
- 14- Friese A, Schulz J, Zimmermann K, Tenhagen BA, Fetsch A, Hartung J, Rösler U. Occurrence of Livestock-Associated Methicillin-Resistant Staphylococcus aureus in Turkey and Broiler Barns and Contamination of Air and Soil Surfaces in Their Vicinity. *Applied and Environmental Microbiology*. 2013;79(8):2759–2766. <https://doi.org/10.1128/aem.03939-12>
- 15- Wang X, Liu Q, Zhang H, Li X, Huang W, Fu Q, Li M. Molecular characteristics of community-associated Staphylococcus aureus isolates from pediatric patients with bloodstream infections between 2012 and 2017 in Shanghai, China. *Frontiers in Microbiology*. 2018; 9(1): 23-29. <https://doi.org/10.3389/fmicb.2018.01211>
- 16- Hado H, Assafi M. Molecular fingerprinting of methicillin resistant Staphylococcus aureus strains isolated from human and poultry in Duhok, Iraq. *Iraqi Journal of Veterinary Sciences*. 2021;35(1):99–103. <https://doi.org/10.33899/ijvs.2020.126375.1310>
- 17- Okonko IO, Nkang AO, Eyarefe OD, Abubakar MJ, Ojezele MO, Amusan TA. Incidence of multi-drug resistant (MDR) organisms in some poultry feeds sold in Calabar metropolis, Nigeria. *British Journal of Pharmacology and Toxicology*. 2010;1(1):15-18. <https://www.airitilibrary.com/Article/Detail/20442467-201007-201008160045-201008160045-15-28>
- 18- Begum SMS, Azeez A, Kavitha K. Isolation, Identification and Characterization of Poultry Associated Bacterial Pathogens. *Uttar Pradesh journal of zoology*. 2013; 44(14):35–41. <https://doi.org/10.56557/upjzo/2023/v44i143556>
- 19- Hossain MT, Siddique MP, Hossain FMA, Zinnah MA, Hossain MM, Alam MK. Isolation, identification, toxin profile and antibiogram of E. coli isolated from broilers and layers in Mymensingh district of Bangladesh. *Bangladesh Journal of Veterinary Medicine*. 2008; 6(1):1-5. <https://www.banglajol.info/index.php/BJVM/article/view/1330>
- 20- Eja ME, Samuel-Udoekong N, Ikpeme EM, Enyi-Idoh KH, Lennox JA, Etim KD. Antibiogram studies and extended spectrum beta-lactamase activity profile of Salmonella-like species isolated from poultry soil of the University of Uyo, Nigeria. *Malaysian Journal of Microbiology*.2012;8(4):280-284. <https://doi.org/10.21161/MJM.44212>
- 21- Igbinosa IH. Antibiogram profiling and pathogenic status of Aeromonas species recovered from chicken. *Saudi Journal of Biological Sciences*.2014; 21(1):481-485. <https://doi.org/10.1016/j.sjbs.2014.06.003>
- 22- Khan MS, Akhtar N, Haque ME, Barua A, Chowdhury T, Mullick R. Isolation and identification of nonplasmid multidrug resistant E. coli from poultry wastes in Chittagong region, Bangladesh. *Journal of Bacteriology and Parasitology*.2014; 5(1):1-7. <https://doi.org/10.4172/2155-9597.1000182>
- 23- Laban SE, Moustafa GZ, Anwer W, Badawy EM. Microbial load of poultry by-products following rendering process. *Global*. 2014;12(6):756-759. [https://idosi.org/gv/gv12\(6\)14/4.pdf](https://idosi.org/gv/gv12(6)14/4.pdf)
- 24- Cuny C, Friedrich A, Kozzytska S. Emergence of methicillin-resistant Staphylococcus aureus (MRSA) in different animal species. *International Journal of Medical Microbiology*. 2010; 300(3): 109–117. <https://doi.org/10.1016/j.ijmm.2009.11.002>
- 25- Ferreira JP, Anderson KL, Correa MT. Transmission of MRSA between companion animals and infected human patients presenting to outpatient medical care facilities, PLoS One. 2011;6(11):26978-26991. <https://doi.org/10.1371/journal.pone.0026978>
- 26- Suleiman A, Zaria LT, Grema HA, Ahmadu P. Antimicrobial resistant coagulase positive Staphylococcus aureus from chickens in Maiduguri, Nigeria. *Sokoto Journal of Veterinary Sciences*. 2013; 11 (1): 51-55. <https://doi.org/10.4314/sokjvs.v11i1.8>
- 27- Assafi MS, Hado HA, Abdulrahman IS. Detection of methicillin-resistant Staphylococcus aureus in broiler and broilers farm workers in Duhok, Iraq by using conventional and PCR techniques. *Iraqi Journal of Veterinary Sciences*. 2020;34(1):15–22. <https://doi.org/10.33899/ijvs.2019.125757.1145>
- 28- Sonola VS, Katakweba A, Misinzo G, Matee MI. Multiplex PCR detection of antibiotic resistance and virulence genes in multidrug-resistant Staphylococcus



- aureus isolated from chickens, humans, rodents, and soil in Northern Tanzania. *German Journal of Microbiology*. 2023;3(2):1–11. Internet Archive. <https://doi.org/10.51585/gjm.2023.2.0024>
- 29- Abd El- Tawab A, Alekhnawy KI, Talaie A. Virulence and resistant genes detection of *Staphylococcus aureus* associated with arthritis in chickens. *Benha Veterinary Medical Journal*. 2018; 35(2):96–106. <https://doi.org/10.21608/bvmj.2018.95986>
- 30- Cheung GYC, Bae JS, Otto M. Pathogenicity and virulence of *Staphylococcus aureus*. *Virulence*. 2021;12(1):547–569. <https://doi.org/10.1080/21505594.2021.1878688>
- 31- Sheet OH, Hussein SA, Al-Chalaby AY. Detection of methicillin-resistant *Staphylococcus aureus* from broiler carcasses in Mosul city. *Iraqi Journal of Veterinary Sciences*. 2021;35(3):489–493. <https://doi.org/10.33899/ijvs.2020.127052.1451>
- 32- Markey B, Leonard F, Archambault M, Cullinane A, Maguire D. *Clinical Veterinary Microbiology*. 2nd Edition, Dublin: Mosby Ltd.2013
- 33- Rahman MM, Amin KB, Rahman SMM, Khair A, Rahman M, Hossain A. Investigation of methicillin-resistant *Staphylococcus aureus* among clinical isolates from humans and animals by culture methods and multiplex PCR. *BMC Veterinary Research*. 2018; 14(1):1-7. <http://dx.doi.org/10.1186/s12917-018-1611-0>
- 34- AL-Salihi SS, Karim GF, Al-Bayati AMS, Obaid HM. Prevalence of Methicillin-Resistant and Methicillin Sensitive *Staphylococcus aureus* Nasal Carriage and their Antibiotic Resistant Patterns in Kirkuk City, Iraq. *Journal of Pure and Applied Microbiology*. 2023;17(1):329–337. <https://doi.org/10.22207/jpam.17.1.22>
- 35- Kanaan MHG. Antibacterial effect of ozonated water against methicillin-resistant *Staphylococcus aureus* contaminating chicken meat in Wasit Province, Iraq. *Veterinary World*. 2018;11(10):1445–1453. <https://doi.org/10.14202/vetworld.2018.1445-1453>
- 36- Aziz SN, Huseein KA, Mohammed AA, Najm TA. Isolation and Identification of Methicillin-Resistant *Staphylococcus aureus* (MRSA) From Different Sources of Food in Basra Province / Iraq. *Egyptian Journal of Medical Microbiology*.2024 33(4): 33-40. <https://doi.org/10.21608/ejmm.2024.311434.1303>
- 37- Richter A, Sting R, Popp C, Rau J, Tenhagen BA, Guerra B, Hafez HM, Fetsch A. Prevalence of types of methicillin-resistant *Staphylococcus aureus* in turkey flocks and personnel attending the animals. *Epidemiology and Infection*. 2012; 140(12):2223–2232. <https://doi.org/10.1017/s095026881200009x>
- 38- Tang Y, Larsen J, Kjeldgaard J, Andersen PS, Skov R, Ingmer H. Methicillin-resistant and -susceptible *Staphylococcus aureus* from retail meat in Denmark. *International Journal of Food Microbiology*. 2017;249(1):72–76. <https://doi.org/10.1016/j.ijfoodmicro.2017.03.001>
- 39- Suk S, Nam L, Park H, Lee H, Choi H, Jung M, Santos SI, Mato R, de Lencastre H, Tomasz AS, Lee J, Kim Y, Sang S, Wee S. Prevalence and characterization of methicillin-resistant *Staphylococcus aureus* in raw meat in Korea. *Journal of microbiology and biotechnology*.2010;20(4):775-778. <https://doi.org/10.4014/jmb.0912.12022>
- 40- Abd El-Ghany WA. *Staphylococcus aureus* in poultry, with special emphasis on methicillin-resistant strain infection: A comprehensive review from one health perspective. *International Journal of One Health*. 2021;1(1):257–267. <https://doi.org/10.14202/ijoh.2021.257-267>
- 41- Hussein NR, Alyas A, Majeed M, Assafi MS. Prevalence rate and prevalent genotypes of CA-MRSA in Kurdistan region: First report from Iraq. *Int J Pure Appl Sci Technol*.2015; 27(1):44-49. <https://doi.org/10.17795/iji-35375>
- 42- Lee JH, Jeong JM, Park YH, Choi SS, Kim YH, Chae JS. Evaluation of the methicillin-resistant *Staphylococcus aureus* (MRSA)-screen latex agglutination test for detection of MRSA of animal origin. *Journal of Clinical Microbiology*. 2004;42(6):2780–2782; <https://doi.org/10.1128/JCM.42.6.2780-2782.2004>
- 43- Paterson GK, Harrison EM, Holmes MA. The emergence of mecC methicillin-resistant *Staphylococcus aureus*. *Trends Microbiology*. 2014;22(1):42–47. <https://doi.org/10.1016/j.tim.2013.11.003>
- 44- Wertheim HF, Melles DC, Vos MC, van leeuwen W, Van Belkum A, Verbrugh HA, Nouwen JL. The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infectious Diseases*.2005;5(12):751-762. [https://doi.org/10.1016/S1473-3099\(05\)70295-4](https://doi.org/10.1016/S1473-3099(05)70295-4)
- 45- Miro RF, Jwher DhM. The Relation Between Biosecurity Measures and the Level of *Staphylococcus aureus* Contamination in Poultry Farms in Northern Iraq. *Journal of Applied Veterinary Sciences*. 2025;10(2):0-0. https://javs.journals.ekb.eg/article_414262_13fb9a517663b575ccd1390751a05b36.pdf