



Molecular Diagnostic Study of Intestinal Protozoa in Wild Pigeons in Diwaniyah Province

Hind Qasim Neamah Al Qasimi¹ Khaled Thamer Matter alshaebani²

Department of Life Sciences, College of Education, University of Al-Qadisyah, Diwaniyah, Iraq

Submitted: May 16, 2025

Revised: June 1, 2025

Accepted: June 04, 2025

Abstract Parasitic infestations in wildlife are significant biomarker of the present state of ecosystems, and studying these infections is essential for better understanding natural epidemiological hotspots. Animals infected with parasites are therefore susceptible to pathogens of medical and veterinary importance, wild birds have an important role in transmitting and spreading many zoonotic diseases, which has a great impact on the health of humans and birds. In this study, samples were collected from the intestinal contents of (50) wild pigeons from different areas in the city of Diwaniyah during the period extending from the beginning of September 2024 to the end of March 2025. The birds were dissected and samples were taken from the intestinal contents to investigate intestinal protozoa. The results of the microscopic examination showed the infection of 14 infected pigeons were frozen and preserved for polymerase chain reaction (PCR) testing. The results revealed two genera of intestinal protozoa, *Cryptosporidium* sp. and *Isospora* sp., with infection rates of 85.71% and 14.28%, respectively.

Keywords: *Cryptosporidium* spp., *Isospora* spp., intestinal protozoa

©Authors, 2025, College of Veterinary Medicine, University of Al-Qadisyah. This is an open access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Birds are among the most important animals in the environment as they are essential for maintaining ecological balance and also serve as seed dispersers, pollinators, and prey regulators. They are considered vital indicators for environmental conservation. They are animals known for their high sensitivity to habitat changes. The extinction of wild species leads to the deterioration of natural environments, affecting not only animals but also plants and microbes, and indirectly contributes to the emergence and spread of parasitic diseases. Parasites may be a major player in maintaining this biodiversity by maintaining diversity in ecological communities^[1]. Wild and domestic birds also play an important role in transmitting and spreading many zoonotic diseases (parasitic, bacterial, viral and fungal), which has a significant impact on the health of humans and birds^[2]. Wild birds are often responsible for contaminating water sources with zoonotic diseases due to their presence near water sources and in urban areas and their ability to transmit intestinal pathogens over long distances^[3]. Parasitic infection in wildlife is an important biomarker of the current state of ecosystems and the study of these infections is essential for a better understanding of natural foci by epidemiology. Animals infected with parasites are therefore susceptible to pathogens of medical and veterinary significance^[4]. The digestive

system is more susceptible to infection by various types of parasites than other organs of the body due to the entry of water or food containing the infective stage of the parasite through the digestive system^[5]. The species of *Cryptosporidium* sp. and *Coccidia* parasites that cause coccidiosis, including species of the genus *Isospora* sp., are considered common intestinal protozoa in birds^[6]. The species of *Cryptosporidium* sp. cause cryptosporidiosis. It is considered a parasitic disease of animal origin that infects multiple hosts, including birds^[7]. The oocyst is considered the infectious stage, as it is infectious when it is excreted from the host's body. It is characterized by its tolerance to environmental conditions and standard water treatments^[8]. Wild birds infected with the parasite also pose a threat to public health due to the nature of their life in the wild, which leads to the difficulty of controlling their infection with *Cryptosporidium*, which increases the possibility of this disease being transmitted to humans and pets. It is also necessary to monitor infection in wild birds, its methods of transmission, and its prevention^[9]. The genus *Isospora* sp. is considered one of the causes of coccidiosis. The oocysts contain two sporocysts, and each sporocyst contains four spores^[10]. Many studies have shown that birds are infected in different organs of the bird's body. In wild birds, it was found in the intestines, liver, and spleen. The stages of *Isospora* sp. in the tissues located

outside the intestines are often accompanied by lymphocytic inflammation of varying severity [11].

Materials and methods

Ethical Approval

The procedures for the current study were approved by the Department of Life Sciences, College of Education. In this study, samples of the intestinal contents of (50) wild pigeons were collected from different areas in the city of Diwaniyah during the period extending from the beginning of September 2024 to the end of March 2025. The results of microscopic examination using traditional methods showed that 14 wild pigeons were infected. These were examined using the conventional polymerase chain reaction technique according to the method of [12]. To detect *Cryptosporidium* sp. and *Isospora* sp. parasites, DNA was extracted using the Presto™ Stool DNA Extraction Kit from Geneaid/Taiwan, which was

provided by Scientific Researcher. Co. Ltd. The extraction was carried out according to the steps in the company's instruction manual. Nano-drop spectrophotometer was used to measure the concentration and purity of DNA, and primers specific to the Small-subunit 18 Sr RNA gene were used to detect *Cryptosporidium* sp. and *Isospora* sp. using PCR technology, as shown in Table (1). These primers were prepared by Scientific Researcher. Co. Ltd, Iraq, where it was designed using the NCBI-Genbank gene bank and the Primer 3 Plus design program. A Nano-drop spectrophotometer was then used to measure the concentration and purity of the extracted DNA. The polymerase chain reaction mixture was prepared in tubes and then transferred to a BIO RAD T 100 Thermocycler for thermal cycling. The PCR product was analyzed by Agarose gel electrophoresis.

Table (1) shows the primers used in this study.

Primer	Sequence 5'- 3'		Product size
<i>Cryptosporidium</i> sp	F	AGACGGTAGGGTATTGGCCT	784bp
	R	TACGAATGCCCCCAACTGTC	
<i>Isospora</i> sp	F	ATTTCTGTCGGTGCATCGG	423bp

Results

PCR results of the intestinal contents of 14 wild pigeons tested showed infection with *Cryptosporidium* sp. (12/14) birds, representing 85.71%, and *Isospora* sp. (2/14) birds, representing 14.28%. Agarose gel electrophoresis of the DNA amplification results from the tested samples for *Cryptosporidium* sp. PCR using the small subunit ribosomal RNA gene primer revealed DNA bands of approximately 784 bp, as shown in Figure (1). Electrophoresis on agarose gel of the DNA amplification product of the samples examined to diagnose the *Isospora* sp parasite using the primer specific to the Small subunit ribosomal RNA gene showed DNA bands of approximately (423 bp) as in Figure (2).

Figure (1): Agarose electrophoresis image showing the analysis of the polymerase chain reaction product of the small subunit ribosomal RNA in *Cryptosporidium* from bird samples. Where: M represents the index for the PCR product (2000-100bp) and the lines (S2, S4, S7, S8 and S9) are some positive samples of *Cryptosporidium* sp at (784 bp).



Figure (2): Agarose electrophoresis image showing the analysis of the polymerase chain reaction product of the small subunit ribosomal RNA in *Isospora* sp. from bird samples. Where: M represents the index for the PCR product (2000-100bp) and the line (S1,S4) for the positive *Isospora* sp samples when the product appears around (423bp).



Discussion

Wild pigeons are found in large numbers around us and may be infected with or harbor numerous types of parasites that can be transmitted to humans or other animals. [13] The use of PCR-based diagnostic methods has increased significantly in the field of parasitology in recent decades. [14] This technique has had a significant impact on the progress achieved in the field of diagnosing, characterizing, tracking, and investigating the evolution, epidemiology, and biology of protozoan parasites. [15] The conventional PCR technique was used in this study, and the infection rate with *Cryptosporidium* spp. in wild pigeons was recorded at 85.71%, which is consistent with what was recorded by [16]. A 90% infection rate of pigeons with the parasite was recorded in a study of primary infection in clinically ill pigeons in Assiut, Egypt. The rate in our current study was close to the rate recorded by [17], who recorded 62% infection in pigeons in his study of Cryptosporidiosis in domestic pigeons in Iran. The infection rate in the current study was higher than that recorded by [18], as the infection rate of wild birds in their study was 25.71% using Nested PCR technology. This may be due to the fact that the samples examined in their study were larger in number, and also the samples in our current study were previously diagnosed positive upon microscopic examination to diagnose the parasite *Cryptosporidium* spp., so it is considered a selective sample and not a random one, several studies have confirmed the prevalence of the *Cryptosporidium* sp. parasite in pigeons worldwide, and

molecular methods have yielded higher rates of infection compared to traditional methods. The highest infection rates were observed in Iraq [19]. The results of the current study also recorded that wild pigeons were infected with the *Isospora* sp. parasite, which was recorded at 14.28%, which is close to what was recorded by [20], who recorded a 19.4% infection rate in pigeons when studying intestinal parasites in pigeons in Venezuela. Several studies have confirmed bird infections with the *Isospora* sp. [21] recorded a 2% infection rate in ostriches when studying the isolation of intestinal parasites from ostriches from central Iraq. [22] also recorded a 2.91% infection rate in domestic birds with the *Isospora* sp. parasite in Bangladesh. [23] He also recorded that wild birds in Europe were infected with 5.8% of the parasite. [24] He also mentioned that wild birds in Britain and Ireland were infected with 32% of the parasite when he studied the prevalence of parasites in 13 families of wild passerine birds. He mentioned that the severity of infection was higher in birds that used a wider range of habitats and increased in areas that contained feeders. The infection of birds in these studies was either higher or lower than what was recorded in our current study. The reason may be due to the difference in the types and numbers of birds studied and the climatic conditions.

Conflict of interest

No conflict of interest is found for the present study.

Funding source

This research had no specific fund; however, it was self-funded by the authors

References

- 1- Oliveira, M. D. S. (2021). Taxonomia e Ecologia de coccídios: *Isospora* spp. e *Eimeria* spp. de aves Silvestres do Brasil e de Portugal.
- 2- Malik, Y. S., Milton, A. A. P., Ghatak, S., & Ghosh, S. (2021). Role of birds in transmitting zoonotic pathogens. Springer Singapore, 276pp.
- 3- Egan, S., Barbosa, A. D., Feng, Y., Xiao, L., & Ryan, U. (2024). The risk of wild birds contaminating source water with zoonotic *Cryptosporidium* and *Giardia* is probably overestimated. *The Science of the total environment*, 912, 169032. <https://doi.org/10.1016/j.scitotenv.2023.169032>
- 4- Lignon, J. S., Pinto, D. M., Dos Santos, T. S., Meireles, G. R., da Silveira, C. G., Bohm, B. C., ... & Bruhn, F. R. P. (2025). Survey of parasitic fauna data from wild animals through coproparasitological diagnosis in Southern Brazil. *BMC Veterinary Research*, 21(1), 7.
- 5- De Oliveira, F. C. R., Gallo, S. S. M., Elizeu, T. K. S., & Ederli, N. B. (2022). Molecular and phylogenetic characterization of *Cryptosporidium* species in the saffron finch *Sicalis flaveola*. *BMC Veterinary Research*, 18(1), 449.
- 6- Gimah, M., Oluwasemilore, J., & Iseal, S. (2025). Impact of Gastrointestinal Parasites on the Health of Captive Wild Animals.
- 7- Essam, A., Abbas, I., Elmishmishy, B., & Elwafa, S. A. (2024). Cryptosporidium infection in ducks and pigeons collected from live bird markets in two Nile Delta governorates (Dakahlia and Gharbia), Egypt. *Mansoura Veterinary Medical Journal*, 25(4), 4.
- 8- Cacciò, S. M., & Widmer, G. (Eds.). (2013). *Cryptosporidium: parasite and disease*. Springer Science & Business Media.
- 9- Wang, Y., Zhang, K., Chen, Y., Li, X., & Zhang, L. (2021). Cryptosporidium and cryptosporidiosis in wild birds: A One Health perspective. *Parasitology Research*, 120(9), 3035-3044.
- 10- Yang, R., Brice, B., Oskam, C., Zhang, Y., Brigg, F., Berryman, D., & Ryan, U. (2017). Characterization of

- two complete *Isospora* mitochondrial genomes from passerine birds: *Isospora serinuse* in a domestic canary and *Isospora manorinae* in a yellow-throated miner. *Veterinary parasitology*, 237, 137-142.
- 11- Kubiski, S. V., Witte, C., Burchell, J. A., Conradson, D., Zmuda, A., Barbon, A. R., ... & Rideout, B. A. (2022). Mitochondrial gene diversity and host specificity of *Isospora* in passerine birds. *Frontiers in Veterinary Science*, 9, 847030.
- 12- Marhoon, I. A., & Al-maeahi, A. M. (2020). Survey of *Isospora belli* infection in some different hosts: birds and mammals. *Indian J Public Health Res Dev*, 11(4), 1213-1218.
- 13- Altamimi, M. K., & Al-Zubaidi, M. T. (2020). *Cryptosporidium* spp.: conventional and molecular detection in wild pigeons (*Columba livia*) in Babylon province, Iraq. *Plant Archives*, 20(2), 1544-1548.
- 14- Baltrušis, P., & Höglund, J. (2023). Digital PCR: modern solution to parasite diagnostics and population trait genetics. *Parasites & Vectors*, 16(1), 143.
- 15- McConkey, G. A. (2024). Rising stars in parasite and host: 2023. *Frontiers in Cellular and Infection Microbiology*, 14, 1509107.
- 16 - ABD EL-SALAMA, A. M., DYAB, A. K., ARAFA, M. I., NASR, A. A., & ABD-ELRAHMAN, S. A. L. W. A. (2025). PROTOZOAL AND RICKETTSIAL INFECTIONS AMONG CLINICALLY ILL PIGEONS IN ASSIUT, EGYPT: PREVALENCE, BIODIVERSITY, AND POTENTIAL PUBLIC RISK. *Assiut Veterinary Medical Journal*, 71(184), 672-683.
- 17- Khordadmehr, M. , Sazmand, A. , Oryan, A. , Almasi, P. , Ranjbar, V. , Hemphill, A. and Otranto, D. (2024). Respiratory and intestinal zoonotic cryptosporidiosis in symptomatic domestic pigeons (*Columba livia domestica*) in Tabriz, Iran. *Iranian Journal of Veterinary Research*, 25(4), 326-332. doi: 10.22099/ijvr.2024.49762.7335
- 18- Al –zubaidi, Niran abed obada & Al Khafaji, Alaa Mohammed Abdulrazaq (2025) Molecular detection and phylogenetic of *Cryptosporidium* spp. of birds in the city of Al-Diwaniyah, Iraq. *Al-Qadisiyah J. Vet. Med. Sci.* 2025; 24 (1): 104-109.
- 19- Devulder, A., Polack, B., & Mammeri, M. (2025). A global review of *Cryptosporidium* spp. in pigeons in the context of the One-Health concept. *Revue d'Elevage et de Médecine Vétérinaire des Pays Tropicaux*, 78, 1-13.
- 20- Cazorla Perfetti, D., & Morales Moreno, P. (2019). Intestinal parasites in feral populations of domestic pigeons (*Columba livia domestica*) in Coro, Falcon State, Venezuela.
- 21- Makawi, Z. A., & Hussein, H. H. (2022). Isolation and identification of intestinal parasites in ostriches *Struthio camelus* Linnaeus, 1758 from Middle of Iraq. *Biochemical and Cellular Archives*, 22(1), 3177-3181.
- 22 - Bayzid, M., Yasir Hasib, F. M., Hasan, T., Hassan, M. M., Masduzzaman, M., Hossain, M. A., & Alim, M. A. (2023). Prevalence of helminth and protozoan infections in pet birds of Chattogram, Bangladesh. *Veterinary Medicine and Science*, 9(1), 548-556.
- 23- Keckeisen, C., Šujanová, A., Himmel, T., Matt, J., Nedorost, N., Chagas, C. R., ... & Harl, J. (2024). *Isospora* and *Lankesterella* Parasites (Eimeriidae, Apicomplexa) of Passeriform Birds in Europe: Infection Rates, Phylogeny, and Pathogenicity. *Pathogens*, 13(4), 337.
- 24- Parsa, F. R., Bayley, S., Bell, F., Dodd, S., Morris, R., Roberts, J., ... & Dunn, J. C. (2023). Epidemiology of protozoan and helminthic parasites in wild passerine birds of Britain and Ireland. *Parasitology*, 150(3), 297-310.